

<https://doi.org/10.17221/96/2021-PSE>

The overview of existing knowledge on medical cannabis plants growing

MATĚJ MALÍK, JIŘÍ VELECHOVSKÝ, PAVEL TLUSTOŠ*

Department of Agroenvironmental Chemistry and Plant Nutrition, Faculty of Agrobiological Sciences, Food and Natural Resources, Czech University of Life Sciences Prague, Prague 6 – Suchbátka, Czech Republic

**Corresponding author: tlustos@af.czu.cz*

Citation: Malík M., Velechovský J., Tlustoš P. (2021): The overview of existing knowledge on medical cannabis plants growing. *Plant Soil Environ.*, 67: 425–442.

Abstract: The use of cannabis for medicinal purposes dates back well before the era of modern medicine, but in recent years research into the use of medical cannabis in the medical and pharmaceutical sciences has grown significantly. In European countries, most cannabis plants have been and still are grown for industrial purposes. For this reason, hemp cultivation technology is relatively well researched, while little is known about the key factors affecting cannabis cultivation for medical purposes. The active substances of cannabis plant targeted by this review are called phytocannabinoids. The biosynthesis of phytocannabinoids is relatively well understood, but the specific environmental factors that influence the type and number of phytocannabinoids have been much less studied. Indoor or greenhouse cultivation, which uses automated lighting, ventilation, irrigation systems and complex plant nutrition has become much more sophisticated and appears to be the most effective method for producing medical cannabis. There are many different cultivation systems for cannabis plants, but one of the essential elements of the process is an optimal plant nutrition and selection of fertilisers to achieve it. This review summarises the existing knowledge about phytocannabinoid biosynthesis and the conditions suitable for growing plants as sources of medical cannabis. This review also attempts to delineate how nutrient type and bioavailability influences the synthesis and accumulation of specific phytocannabinoids based on contemporary knowledge of the topic.

Keywords: *Cannabis sativa* L.; tetrahydrocannabinol; cannabidiol; chemical profile; growing conditions

Cannabis is one of the earliest of domesticated crops. According to Chinese historical records and archaeological findings, its cultivation and utilisation can be traced back to 3 000 to 4 000 years BCE (Yu 1987, Jiang et al. 2006). The first use of cannabis for therapeutic purposes, directly evidenced by the finding of the stable cannabis compound, Δ^6 -tetrahydrocannabinol (Δ^6 -THC), has been dated to around 400 CE in a carbonised material discovered in a tomb at Beit Shemesh near Jerusalem (Zlas et al. 1993). Recent years have seen a boom in research on medical cannabis in the biomedical and pharmaceutical sectors. The applicability and acceptability of medical cannabis is expanding, as seen by the growing number of countries that allow its use for specific therapeutic indications (Shelef et al. 2011, Troutt and

Didonato 2015, Balneaves and Alraja 2019). The number of active phytocannabinoids under investigation continues to increase and their effects on a variety of diseases such as chronic pain (Lynch and Campbell 2011, Portenoy et al. 2012, Wilsey et al. 2013), nausea and vomiting (Lane et al. 1991, Duran et al. 2010), spasticity (Pooyania et al. 2010, Corey-Bloom et al. 2012), depression (Wade et al. 2004, Selvarajah et al. 2010, Portenoy et al. 2012), glaucoma (Järvinen et al. 2002), inflammatory bowel disease (Ravikoff Allegretti et al. 2013), psychosis, motor and non-motor symptoms of Parkinson disease (Lotan et al. 2014), anxiety and sleep disorder (Russo et al. 2007, Bonn-Miller et al. 2014, Babson et al. 2017) are being studied (Doyle and Spence 1995, Järvinen et al. 2002, Lynch and Campbell 2011, Grotenhermen and

Muller-Vahl 2012, Ravikoff Allegretti et al. 2013, Lotan et al. 2014). Nearly 150 different phytocannabinoid compounds are currently known (Hanuš et al. 2016).

TAXONOMY

History

The genetic plasticity of cannabis makes it difficult to catalog, and there is still a debate about its proper botanical classification. Linnaeus (1753) described *Cannabis sativa* as a single species. Based on comparative analyses of the psychoactive effects, leaf size, shape and structure of Indian and European varieties, Jean-Baptiste Lamarck (1786) classified the Indian cultivars as a separate species, *Cannabis indica*. At the beginning of the 20th century, the Russian botanist Janischevsky (1924) found that the local Russian plants possessed different characteristics from both *C. sativa* and *C. indica* yet still belonged to the cannabis taxon. These small, wild-growing, auto-flowering plants have been classified as a separate species named *Cannabis ruderalis* (Figure 1).

Current nomenclature

Small and Cronquist (1976) utilised a biphasic approach combining morphological and chemical characteristics to divide the *Cannabis* genus into the following four groups:




Figure 1. Species of cannabis (Hartsel et al. 2016)

1. *Cannabis sativa* L. subsp. *sativa* var. *sativa*,
2. *Cannabis sativa* L. subsp. *sativa* var. *spontanea* Vavilov,
3. *Cannabis sativa* L. subsp. *indica* Small & Cronquist var. *indica* (Lam) Wehmer,
4. *Cannabis sativa* L. subsp. *indica* Small & Cronquist var. *kafiristanica* (Vavilov) Small & Cronquist (Figure 2).

Hillig (2005) concluded from his genomic study of the classification of *C. sativa* that none of the previous taxonomic concepts sufficiently defined the *sativa* and *indica* genes. He analysed different genotypes from various geographical origins and was therefore inclined to a multi-species classification including *C. sativa*, *C. indica* and *C. ruderalis*. Small (2015) has recently proposed two possible cannabis taxonomic classifications. The first is consistent with an earlier division (Small and Cronquist 1976) and is in accordance with the International Code of Nomenclature for Algae, Fungi, and Plants (McNeill et al. 2012). The second, for domesticated cannabis, follows the guidelines of the International Code of Nomenclature for Cultivated Plants (Brickell et al. 2009):

Non-narcotic plants, domesticated for stem fiber and/or oilseeds in West Asia and Europe. Low Δ^9 -tetrahydrocannabinol (THC) content and high cannabidiol (CBD) content (Hillig and Mahlberg (2004) *Cannabis sativa* "hemp biotype").

Non-narcotic plants, domesticated for stem fiber and/or oilseeds in East Asia, mainly China. From low to moderate THC content and high CBD content (Hillig and Mahlberg (2004) *Cannabis indica* "hemp biotype").

Psychoactive plants, domesticated in Southern and Central Asia. High THC content and low or absent CBD content (Hillig and Mahlberg (2004) *Cannabis indica* "narrow-leaflet drug (NLD) biotype").

Psychoactive plants, domesticated in Southern Asia (Afghanistan and neighboring countries). From moderate to high THC and CBD content (Hillig and Mahlberg (2004) *Cannabis indica* "wide-leaflet drug (WLD) biotype").

In addition, two hybrid classes have also been generated:

5. Non-narcotic plants, hybrid cultivars between two fiber (hemp) groups (1 and 2).
6. Psychoactive plants, hybrid cultivars between two narcotic groups (3 and 4).

Hillig and Mahlberg (2004) analysed the content of cannabinoids in various cannabis plants and based on geographical origins, morphological features and

<https://doi.org/10.17221/96/2021-PSE>




Figure 2. Cannabis chemotypes (Small and Cronquist 1976). THC – Δ^9 -tetrahydrocannabinol

the supposed purpose of cultivation assigned them to the intraspecific taxa (biotypes):

Cannabis sativa "hemp biotype" – 62 plants were analysed, ranges of the dry-weight percentages of THC were measured 0.1–11.5% and CBD were measured 0.0–13.6%.

Cannabis indica "hemp biotype" – 45 plants were analysed, ranges of the dry-weight percentages of THC were measured 0.1–9.3% and CBD were measured 0.0–8.5%.

Cannabis indica "narrow-leaflet drug (NLD) biotype" – 68 plants were analysed, ranges of the dry-weight percentages of THC were measured 1.4–12.4% and CBD were measured 0.0–0.1%.

Cannabis indica "wide-leaflet drug (WLD) biotype" – 40 plants were analysed, ranges of the dry-weight percentages of THC were measured 0.1–14.7% and CBD were measured 0.0–11.0%.

All cannabis species successfully cross and produce fertile hybrids (Beutler and Marderosian 1978). *Indica* and *sativa* plants have also been found to differ in terpene and cannabinoid profiles. Thus, these chemotaxonomic markers are a promising tool for screening hybrids (Hillig 2004, Hillig and Mahlberg 2004, Fishedick et al. 2010, Elzinga et al. 2015). Zhang et al. (2018) are recommending that Cannabis should be recognised as a monotypic species typified by *Cannabis sativa* L., containing three subspecies: subsp. *sativa*, subsp. *indica*, and subsp. *ruderalis*.

This proposal is based on their study focused on DNA sequence variations of cannabis plants. Also, McPartland (2018) in his work mentions that DNA barcode analysis supports the separation cannabis at a subspecies level and recognising the nomenclature of *C. sativa* subsp. *sativa* and *C. sativa* subsp. *indica*.

BIOSYNTHESIS OF CANNABINOIDS

History

Actual cannabinoid research is based on a number of major discoveries made by Professor Raphael Mechoulam and Professor Yechiel Gaoni. In the 1960's they identified the psychoactive component in *Cannabis sativa*, Δ^9 -tetrahydrocannabinol, determined and described its chemical structure (Gaoni and Mechoulam 1964, Mechoulam and Gaoni 1967) and synthesised it (Mechoulam et al. 1967). Endogenous cannabinoid receptor ligands, called endocannabinoids, were identified in mammalian tissues in the 1990s. The best-known examples are anandamide (Devane et al. 1992) and 2-arachidonoyl-glycerol (Mechoulam et al. 1995). Endocannabinoids are derived from arachidonic acid, and membrane lipids serve as a potential source of this fatty acid (Giuffrida et al. 2001). For this reason, cannabinoids from cannabis are often referred to as phytocannabinoids to differentiate them from endocannabinoids.

Biosynthesis of phytocannabinoids

Phytocannabinoids can be divided into two groups, neutral cannabinoids and cannabinoid acids. Diversification is based on how many carboxyl groups the molecule has, but non-enzymatic decarboxylation can occur during storage and especially at elevated temperatures when cannabis is smoked (Kimura and Okamoto 1970, Shoyama et al. 1970). Phytocannabinoids, prenylated polyketides of mixed biosynthetic origin, are synthesised from fatty acid precursors and isoprenoids. All phytocannabinoid structures contain a monoterpene unit attached to the phenolic ring having the C3 alkylated carbon (Dewick 2002). The alkyl side chain can vary in length from one to five carbons (Figure 3) and n-pentyl is the most abundant (Elsohly and Slade 2005). Phytocannabinoids containing an n-propyl side chain are referred to as cannabivarin. Tetrahydrocannabivarin (THCV), the THC analogue with an n-propyl side chain, often occurs in *C. indica* (Hillig and Mahlberg 2004).

The starting materials for aromatic ring synthesis, including the alkyl on the third carbon (Hanuš et al. 2016), are three molecules of malonyl-CoA and one molecule of hexanoyl-CoA derived from hexanoic (caproic) acid (Dewick 2002). The hexanoyl-CoA acts as a primer for the type III polyketide synthase enzyme, also known as tetraketide synthase (TKS), which also requires the olivetolic acid cyclase enzyme (OAC) catalysing a C2–C7 intramolecular aldol condensation with carboxyl group retention to produce olivetolic acid (Taura et al. 2009, Gagne et al. 2012). These transformations can give rise to by-products such as 4-hydroxy-6-pentylpyran-2-one (PDAL), 4-hydroxy-6-(2-oxoheptyl)pyran-2-one (HTAL) and olivetol. Cannabigerolic acid (CBGA) is further derived from olivetolic acid after alkylation with a monoterpene unit, geranylpyrophosphate, with the participation of geranylpyrophosphate:olivetolate geranyltransferase (GOT) (Figure 4) (Fellermeier and Zenk 1998). Also, the (Z)-isomer of cannabigerolic acid, cannabimerolic acid (CBNRA), is synthesised to a small extent when neryl pyrophosphate is used by




Figure 3. Structure of cannabinoids

the GOT enzyme instead of geranyl pyrophosphate (Taura et al. 1995a). There are three acids that can be formed from CBGA and CBNRA.

Tetrahydrocannabinolic acid (THCA) is produced during the formation of the heterocyclic ring by the THCA synthase enzyme, which can convert CBGA or CBNRA to THCA (Figure 5) (Taura et al. 1995b). However, the low THCA synthase specificity for CBNRA compared to CBGA suggested that THCA was predominantly synthesised from CBGA. The course of this reaction is similar to that of other reactions catalysed by monoterpene cyclases. Most of the cyclases require divalent ions such as Mg^{2+} or Mn^{2+} for their activity, but this is not the case with THCA synthase (Taura 2009). The presence of a carboxyl group in the substrate molecule is essential for the reaction because THCA synthase does not recognize neutral phytocannabinoids such as cannabigerol (CBG) as substrates (Taura et al. 2007a).

The structure of cannabidiolic acid (CBDA) is the result of a pericyclic reaction involving loss of a proton (Figure 6) (Dewick 2002). The modification is catalysed by the intramolecular oxidoreductase, CBDA synthase, which selectively favours the formation of CBDA from CBGA over its (Z)-isomer, CBNRA (Taura et al. 1996). The effects of various metal ions (Mg^{2+} , Mn^{2+} , Zn^{2+} , Ca^{2+} , Co^{2+} and Cu^{2+}) on its activity were investigated, but they did not alter the rate of catalysis. In contrast, the Hg^{2+} ion completely inhibited enzyme activity at a concentration of 2 mmol, and the chelating agent, ethylenediaminetetraacetic acid (EDTA), at concentrations up to 5 mmol showed a low positive effect on enzyme activity. Thus, CBDA synthase does not appear to require metal ions for CBGA oxidocyclization (Taura et al. 1996). CBDA synthase and THCA synthase catalyse the formation of single optical isomers at a purity of greater than 95% (Taura et al. 2007b).

Cannabichromenic acid (CBCA) is derived from CBGA by oxidation and cyclisation by cannabichromenic acid synthase (CBCA synthase) (Figure 7). CBCA is synthesised as a 5:1 enantiomeric mixture, probably because of the partial release of intermediates from the CBCA synthase active site prior to completion of the reaction (Morimoto et al. 1997). Tests of the metal ions, Mg^{2+} , Zn^{2+} , Ca^{2+} and Cu^{2+} , showed that none of them stimulated enzyme activity. Hg^{2+} , however, completely inhibited the reaction at a concentration of 1 mmol. EDTA slightly increased enzyme activity suggesting that the CBCA synthase reaction does not require metal ions (Morimoto et al. 1998).

<https://doi.org/10.17221/96/2021-PSE>




Figure 4. Biosynthesis of phytocannabinoids 1/2. TKS – tetraketide synthase; PDAL – 4-hydroxy-6-pentylpyran-2-one; OAC – olivetolic acid cyclase enzyme; HTAL – 4-hydroxy-6-(2-oxoheptyl)pyran-2-one; GOT – geranylpyrophosphate: olivetolate geranyltransferase




Figure 5. Δ^9 -THCA (tetrahydrocannabinolic acid) synthesis




Figure 6. Cannabidiolic acid (CBDA) synthesis

Croteau (1987) discovered that all terpene cyclases require bivalent cations for their function because these metal ions are able to neutralise the negative charge on the diphosphate groups on the terpene molecules and ionise the allyl diphosphate substrate. Since CBGA does not contain a diphosphate group it is to be expected that CBCA synthase, CBDA synthase and THCA synthase have no requirement for bivalent cations. The most of the cannabinoids present in *C. sativa* can be categorised as Δ^9 -tetrahydrocannabinol (Δ^9 -THC), CBD, CBC, CBG, cannabinol (CBN), cannabicyclol (CBL), cannabielsoin (CBE) and cannabitrinol (CBT) (Turner et al. 1980, Razdan 1986, Ross and Elsohly 1995). Δ^9 -THCA, CBDA and CBCA are also sometimes called primary phytocannabinoids because other phytocannabinoids are generated from these three precursors predominantly by nonenzymatic degradative pathways.

Primary phytocannabinoids can either be decarboxylated to their neutral form (Figure 8) or converted to CBE, CBN, CBT, Δ^8 -tetrahydrocannabinol (Δ^8 -THC) or CBL *via* exposure to light, heat and oxygen (Figure 9). CBD can undergo photooxidation or pyrolysis to form CBE. Δ^9 -THC is converted to the thermodynamically more stable Δ^8 -THC when exposed to heat, or it may be degraded to CBT or CBN in the presence of oxygen (Elsohly and Slade 2005). The presence of CBT and CBN together with high levels of decarboxylated phytocannabinoids, are the chemical indicators of lengthy storage under poor conditions (Shoyama et al. 1970). The degradation rate of primary phytocannabinoids to these

secondary phytocannabinoids increases with higher temperature, higher initial concentrations of primary phytocannabinoids, and with an increase in the inflorescence surface, and thus greater surface exposure to air (Milay et al. 2020). CBC in the presence of light converts to CBL-type phytocannabinoids (Elsohly and Slade 2005). Cannabivarins are generated by the same biosynthetic pathways from cannabigerovarinic acid (CBGVA), a homologous CBGA precursor (Shoyama et al. 1984). The cannabinoid profile in *Cannabis* undergoes rapid changes in the early stages of growth (Potter 2014). CBDA and THCA synthases have very similar catalytic rates ($k_{\text{cat}} = 0.19/\text{s}$ and $0.20/\text{s}$) and affinity ($K_M = 134 \mu\text{mol}$ and $137 \mu\text{mol}$) for cannabigerolic acid (Taura et al. 1995b, 1996). The CBCA synthase, however, shows a lower Michaelis constant ($K_M = 23 \mu\text{mol}$) as well as a higher catalytic rate ($k_{\text{cat}} = 0.04/\text{s}$). In the early stages of cultivation, where CBGA is still present at low concentrations, CBCA synthesis predominates (Morimoto et al. 1998). However, as the CBGA concentration increases over time, the efficacy of THCA and CBDA biosynthesis increases, and these molecules soon outweigh the CBCA concentration. At later stages of growth, CBGA synthesis slows and its relative proportion in the phytocannabinoid profile is gradually reduced (Potter 2014).

CULTIVATION

In European countries, most cannabis is grown for industrial purposes (Zuk-Golaszewska and Golaszewski




Figure 7. Cannabichromenic acid (CBCA) synthesis

<https://doi.org/10.17221/96/2021-PSE>




Figure 8. Decarboxylation of primary phytocannabinoids

2018) such as hemp fibers (Pickering et al. 2007), seeds as a source of oil (Mölleken and Theimer 1997, Kriese et al. 2004) and protein (Patel et al. 1994). For this reason, the procedures for hemp cultivation are well known, while the growth factors affecting cannabis cultivation for medical purposes are poorly understood (Zuk-Golaszewska and Golaszewski 2018).

Indoor or outdoor medical cannabis cultivation?

The conditions under which cannabis plants are grown for drug production is subject to more stringent protocols relating to the content and type of the active phytocannabinoids. Among the factors influencing the composition and yield of phytocannabinoids are the genotype of the plant, the growing conditions, maturity at harvest time, storage and handling (Potter 2014).

It is much more efficient to grow medical cannabis plants in a greenhouse where light, temperature and humidity can be controlled. Until recently, this method of cultivation was used mainly by illegal cannabis growers (Drugs 2009). Outdoor cultivation is less expensive, but the variability of the environment makes it almost impossible to obtain a high-

potency, homogeneous product. Cannabis that is grown outdoors is also at greater risk from pests and plant diseases (Potter 2014). Cannabis entrepreneurs now use sophisticated indoor cultivation methods with automated control of lighting and photoperiod, temperature, ventilation and irrigation, and complex systems for providing nutrients. However, much of the information on indoor cannabis production is still obtained from anecdotal sources (Vanhove et al. 2011). Current data on the influence of photoperiod and even light spectrum allow indoor growers to regulate such aspects as leaf and shoot growth and time of vegetation cycle and thus achieve several growth cycles per year (Farag and Kayser 2015). Three to six harvests per year (six harvests per year is the maximum, and in this case, you have to skip the vegetative phase) can be attained by applying modern controlled growing practices (Leggett 2006).

Hydroponics versus soil

Indoor cannabis cultivation can be accomplished in several ways, but primarily either in soil or in soilless culture using hydroponic media. Hydroponic




Figure 9. Biosynthesis of phytocannabinoids 2/2. CBGA – cannabigerolic acid; CBGVA – cannabigerovarinic acid; CBDA – cannabidiolic acid; CBDVA – cannabidivarinic acid; CBCA – cannabichromenic acid; CBCVA – cannabichromevarinic acid; THCA – tetrahydrocannabinolic acid; THCVA – tetrahydrocannabivarinic acid; CBEA – cannabielsoin acid; CBEVA – cannabielsovarinic acid; CBLA – cannabicyclic acid; CBLVA – cannabicyclvarinic acid; CBTA – cannabitriolic acid; CBNA – cannabinolic acid

cultivation has become increasingly popular among growers. A soilless media such as mineral wool, co-

conut fibers, perlite or expanded clay are used while nutrients provided by solutions are applied directly

<https://doi.org/10.17221/96/2021-PSE>

to the roots (Vanhove et al. 2011). The conventional type of cultivation is in soil with fertilisers applied through irrigation or by mixing with the soil substrates. Potter (2014) found that there was no increase in phytocannabinoid potency or in biomass under hydroponic conditions compared to standard soil cultivation, and the hydroponic system was more complicated and difficult to operate and maintain.

Vegetation cycle of cannabis

Cannabis is a short-day plant, naturally blooming in autumn, and the induction of flowering is regulated by specialised photoreceptor proteins called phytochromes. Therefore, the effect of photoperiod must be taken into an account in indoor cultivation (Halliday and Fankhauser 2003). The vegetative phase lasts from 2–4 weeks after rooting the clones or germinating the seeds (De Backer et al. 2012). The relative humidity in this phase should be from 70% to 80% with a temperature from 21 °C to 28 °C (Chandra et al. 2008). The generative phase is induced by shortening the photoperiod to 12 h light and 12 h dark. The first flowers should appear about one week after the reduction of the light period. The development of stems and leaves gradually slows down and stops after three weeks of this photoperiod, while the flowers continue to develop over the next 8 weeks (De Backer et al. 2012). The monitoring of 200 high THC cannabis varieties showed that the average flowering time with 12 h light period was 57 days, and 88% of the plants flowered between 7 and 9 weeks (Carpentier et al. 2012). The recommended temperatures are similar to the previous phase from 21 °C to 28 °C. However, humidity should be lowered to 40% over the generative phase to reduce the risk of fungal diseases (Vanhove et al. 2011, 2012).

Effect of CO₂ concentration

In order to prevent mold, a dry environment and constant air circulation should be ensured in indoor cannabis growing rooms, either from outdoor ventilation with filters or by indoor fans. It is also recommended to increase the concentration of CO₂ during the light phase of the day (cycle) to improve photosynthesis, plant growth and thus increase biomass yield (Kimball 1983, Wheeler et al. 1996, Chandra et al. 2008, 2011). Elevated CO₂ concentration can improve the assimilation of carbon, thereby accelerate plant growth and potentially improve productivity (Kimball 1983). There is a close correlation between plant yield

and photosynthesis rate because more than 90% of plant dry matter is derived from photosynthetic CO₂ assimilation (Zelitch 1975). However, the improved level of plant photosynthesis and growth appear to be species- and variety-specific (Minorsky 2002).

Wang et al. (2008) investigated the effects of standard (370 ppm) and high (700 ppm) CO₂ concentrations on photosynthesis tolerance to acute heat stress (daily growth temperature was increased by 15 °C every day for 4 h) in cool-season and warm-season of C3 plants. High CO₂ concentration increased the cool-season and warm-season C3 plants tolerance of photosynthesis to acute heat stress. Hamilton et al. (2008) further elaborated the previous idea and concluded that the effects of growth temperature on photosynthetic thermotolerance between C3 and C4 plants are different and affected by the state of acclimatisation of the plants. A high concentration of CO₂ (700 ppm) increases the thermotolerance of C3 plants photosynthesis, except for C3 plants grown at the supra-optimal (5 °C above optimal) growth temperature, then increased CO₂ may provide no advantage or even reduce photosynthesis. On the other hand, increased CO₂ often reduces the photosynthetic thermotolerance of C4 plants at both optimal and supra-optimal growth temperatures.

Chandra et al. (2011) performed experiments directly on cannabis and showed that increasing CO₂ concentration from 390 ppm to 700 ppm increased the rate of photosynthesis in different varieties of *Cannabis sativa* by 38–48% and improved efficiency of water uptake.

Artificial light

To achieve optimal biomass and phytocannabinoid production, artificial lighting must meet certain parameters. These include light intensity in lumens per m² (lux units) and radiation intensity in watts per m² and the wavelength. Wavelength is particularly important because plants require different wavelengths of light during the growth. In the vegetative (roots and shoots) phase, the light should be 420–460 nm which corresponds to blue light, which promotes phototropism and growth hormone production in the plants. In the flowering phase, a red spectrum (600–680 nm) that is well absorbed by chlorophyll is best (Mahlberg and Hemphill 1983). For indoor cannabis cultivation, fluorescent T-5 lighting, metal-halide lamps (MH), high-pressure sodium lamps (HPS) for the growth and light-emitting diodes (LED),

high-pressure sodium lamps (HPS) for the generative phase are most commonly used (Sweet 2016). These lamps differ in the composition of the inside gases, and they produce the light of different wavelengths.

The optimal intensity of illumination. The experiments of Potter and Duncombe (2012) showed positive relationship between the intensity of illumination and amount of biomass harvested. They determined three zones with elevated illumination energy 270, 400 and 600 W/m². Five plants of each variety were placed in each of the three zones at a density of 10 plants/m². In the growth rooms, daily average temperatures were maintained at 25 ± 2 °C. A constant supply of fresh air kept CO₂ concentration in the environment between 350 ppm and 390 ppm. Irradiance levels at the surface of the plant canopy were measured using a hand-held light meter determined the photosynthetically active radiation 80, 120 and 180 W/m² according to variants. Within plants growth, the lamps were kept at a constant distance from the cannabis canopy. The greatest harvest was achieved at 600 W/m² of the illumination intensity. Furthermore, the THC contents in the leaves and inflorescences of the mentioned variants were measured, but no significant increase in the concentration of THC was recorded with an increase of light intensity. Toonen et al. (2006) also reported that plants grown under 600 W lamps achieved higher yields than plants grown under 400 W lamps.

Decreasing tendency of plants to convert light energy into biomass with increasing levels of radiation is probably due to the fact, that plants have a limited ability to use light for photosynthesis. Under low light conditions, plants normally show an initial linear increase in the rate of photosynthesis and thus a tendency to convert light energy into biomass in response to increasing irradiation. However, under brighter conditions, the growth rate slows as chloroplasts become more and more saturated with light (Evans et al. 1993, Ögren and Evans 1993). This has also been proven on cannabis. The rate of increase in photosynthetic activity went down rapidly when irradiation levels rose above 100 W/m² of photosynthetically active radiation. Since 300 W/m² of photosynthetically active radiation, almost no increase in photosynthetic activity has been observed (Lydon et al. 1987).

HPS lamps versus LED. Magagnini et al. (2018) concluded that HPS-lit plants were higher and had a larger amount of dry matter than LED-lit plants. Conversely, plants under LED fixtures contained

higher levels of CBD and THC than under the HPS. Namdar et al. (2019) also found out significant increase in concentration of CBGA in the inflorescences that flowered under LED illumination, with CBGA:THCA ratio of 1:2 as opposed to 1:16 when grown under HPS. Because of the high level of illumination, it was necessary to install a ventilation fan for cooling to the optimum temperature for photosynthesis of 25 °C to 30 °C (Bazzaz et al. 1975). A more efficient alternative is to use banks of LEDs that produce relatively little heat (Bessho and Shimizu 2012). LEDs do not consume much energy, do not require ballasts, and produce only a small amount of heat compared to high intensity discharge lamps. LEDs are compact, have long lives, very good wavelength specificity, relatively cool radiating surfaces, and linear photon output with electrical input current (Massa et al. 2008).

NUTRITION

In the area of plant nutrition for medical cannabis production, there is currently a lack of experimental data in the literature (Caplan et al. 2017a). It is known that the content of cannabinoids in leaves gradually decreases from top to bottom of the hemp plant (Hemphill et al. 1980) and from the literature about hemp cultivation can be deduced that nutrient application can affect the final cannabinoid content of the plants as well as their total yield. This suggests that nutrition could play a similar role for medical cannabis grown under controlled conditions. However, cannabis for hemp production has been selectively bred to produce fiber and is therefore likely to have slightly different nutrient needs than cannabis grown for medicinal purposes. The hemp crop is also grown in the field and not indoors (Hillig and Mahlberg 2004, Van Bakel et al. 2011, Amaducci et al. 2015).

Acceptable forms of individual essential nutrients are divided by Barker and Pilbeam (2015) into two groups according to plant needs, namely macronutrients: nitrogen (NO₃⁻, NH₄⁺), phosphorus (H₂PO₄⁻, HPO₄²⁻), potassium (K⁺), calcium (Ca²⁺), sulfur (SO₄²⁻), magnesium (Mg²⁺), and micronutrients: iron (Fe²⁺, Fe³⁺), chlorine (Cl⁻), manganese (Mn²⁺), zinc (Zn²⁺), copper (Cu⁺, Cu²⁺), boron (H₃BO₃, H₂BO₃⁻), molybdenum (MoO₄²⁻) and nickel (Ni²⁺).

Macronutrients

Nitrogen, phosphorus and potassium (NPK). It is assumed that the nitrogen content in the vegetative parts of

<https://doi.org/10.17221/96/2021-PSE>

the hemp plant positively correlates with the THC content (Haney and Kutscheid 1973). Thus, older leaves contain less THC than younger leaves because they contain less nitrogen. In contrast, high nitrogen levels in applied nitrogen fertilisers reduce the THC content of the hemp leaves (Bócsa et al. 1997). For example, good hemp production requires optimum soil nitrogen levels in the range of 50–200 kg/ha (Vera et al. 2004, Aubin et al. 2015), but these recommendations are not applicable for hydroponic or soil cultivation where studies on indoor cannabis cultivation indicated nitrogen fertilisation should be provided in the range of 190–400 mg N/L. This value has also been reported for nitrogen supplementation of organic, greenhouse-grown tomatoes (Zhai et al. 2009, Surragé et al. 2010).

Hemp growth and an increase in THC content were positively correlated with soil P content (Coffman and Gentner 1977). A negative relationship has been reported for CBD content in leaf tissue relative to available P. Hemp grown on soils depleted of P showed an increased CBD content (Coffman and Gentner 1975). Conversely, phosphorus enhancement did not show any positive effect on THC, CBD, CBN or CBG concentrations in buds from the top of the medical cannabis plants (Bernstein et al. 2019b).

Saloner et al. (2019) investigated response of medical cannabis to different potassium supply in vegetative growing phase. The results show that the response to nutrition is highly dependent on the genotype. Plants in this study were exposed to five different levels of K supply (15–240 ppm). Generally, both cultivars showed increased K concentration in all plant parts with increased K supply. Insufficient K dose for optimal growth and function was the lowest tested supply 15 ppm of K. Also, the highest dose proved excessive and damaging effect to development for one of the two tested genotypes. Similarities proven at both genotypes were in trends of accumulation and uptake. Results demonstrated competition between K and Ca with Mg uptake and no effect on P and N uptake except in the K deficiency range. Potassium supply showed only little effect on micronutrient accumulation in the plant shoot which was similar for both cultivars.

In contrast, no significant effect on hemp biomass and THC was observed in relation to different doses of N and K (Coffman and Gentner 1977). According to Hanuš and Dostálová (1994), various combinations of selected macroelements (N, P, K) in hemp culture can significantly affect the type of phytocannabinoids present and their individual contents. One of a few available sources of scientific literature dealing directly with this issue is the article by Caplan et al. (2017a,

b), who reported a concentration of 389 mg N/L as optimal during the growth phase for maximum yield. The ratio of the basic macroelements (N, P, and K) in the vegetative period was 4:1.3:1.7. After making the calculations for P and K, we obtained values of 126 mg P/L and 165 mg K/L. In the generative phase, 212–261 mg N/L was the optimal amount. A nitrogen concentration of 283 mg N/L gave the maximum yield of inflorescence and biomass, but the concentrations of phytocannabinoids in the dried product was lower. The ratio of N, P, and K in the generative period was set at 2:0.87:3.32. Therefore, an initial concentration of 283 mg N/L, would require 123 mg P/L and 470 mg K/L. The plants tested were propagated from 17 day-old cuttings, which were fertilised with a solution of the indicated concentration for the following 21 days of vegetative growth. Another study has proved sensitivity of phytocannabinoids metabolism to mineral nutrition. The results presented by Bernstein et al. (2019b) show that increased treatment of inorganic NPK increased levels of CBG in flowers by 71% and decreased levels of CBN in flowers by 38% compared to a control treatment. Plants in the control variant were cultivated in potting mixture with fertigation. Concentration of dissolved nutrients in the control variant was as follows: 65 ppm N (1:2 ratio of $\text{NH}_4^+/\text{NO}_3^-$), 17 ppm P (40 ppm P_2O_5), 90 ppm K (108 ppm K_2O). Micronutrients were supplied chelated with EDTA at concentration of 0.4 ppm Fe, 0.2 ppm Mn, and 0.06 ppm Zn.



The rest of macronutrients. The magnesium cation content in soils is relatively mobile and its concentration in plants, especially in leaves, is high because it is a component of chlorophyll. The negative correlation between this metal and copper results from the fact that the radii of their ions are similar and both ions can compete for the same binding sites. The content of Δ^9 -THC and CBD in hemp leaves decreases with increasing Mg concentration in the soil. The Δ^9 -THC content in leaves is positively correlated with the ratio of accessible Ca/Mg in soil. CBD is negatively correlated with available Ca/Zn and Mg/Cu ratios. Positive correlations of magnesium with Δ^8 -THC have been reported with the hypothesis that this nutrient may be cofactor in the enzyme responsible for its production (Coffman and Gentner 1975, Pate 1994, Radosavljevic-Stevanovic et al. 2014).

Micronutrients

Similar results have been seen for micronutrient requirements. Positive correlations of iron with

Δ^8 -THC have been reported with the hypothesis that this nutrient may be cofactor in the enzyme responsible for its production (Pate 1994, Radosavljevic-Stevanovic et al. 2014). CBD content in hemp plants is decreasing with increasing iron concentration (Radosavljevic-Stevanovic et al. 2014). The negative correlation of iron (Fe) and chromium (Cr) with CBD can be explained because the catalase responsible for the decomposition of hydrogen peroxide from the CBDA synthase reaction is a member of the class that contains four heme iron groups. Hydrogen peroxide is strongly sterically hindered upon entry into the heme cavity where the first step of catalysis takes place. Transferring a proton from an oxygen atom to a hydrogen peroxide molecule, and then to a second oxygen atom extends and polarises the O-O bond, which eventually decays heterolytically. The first oxygen atom of the hydrogen peroxide molecule is coordinated with a heme center, which releases water and creates an $\text{O}=\text{Fe}^{\text{(IV)}}\text{-enzyme}^{(+)}$ heme radical. The radical then quickly breaks down by electron transfer, removing the radical electron from the porphyrin ring, which remains unchanged. During the second step, in a similar two-electron transmission reaction, the $\text{O}=\text{Fe}^{\text{(IV)}}\text{-enzyme}^{(+)}$ reacts with a second molecule of hydrogen peroxide to form the parent molecule $\text{Fe}^{\text{(III)}}\text{-enzyme}$, water, and molecular oxygen (Boon et al. 2007, Vlasits et al. 2010).

Proposed reaction mechanism:




Fe-enzyme represents the center of heme iron attached to the rest of the enzyme.

The transition state, $\text{O}=\text{Fe}^{\text{(IV)}}\text{-enzyme}^{(+)}$ is energetically unstable, so these reactions are disadvantageous (Boon et al. 2007, Vlasits et al. 2010). Although chromium is not important for plant growth, its negative correlation with CBD is explained by the fact that Fe and Cr occur together in nature as a complex oxide (Radosavljevic-Stevanovic et al. 2014).

The concentration of CBN and Δ^9 -THC in hemp plants can be influenced by the amount of manganese (Radosavljevic-Stevanovic et al. 2014). A positive correlation of manganese with CBN has been reported (Pate 1994, Radosavljevic-Stevanovic et al. 2014). THCA synthase, which catalyses the oxidative cyclisation of CBGA to THCA, contains a flavin adenine dinucleotide (FAD) prosthetic group that is reduced to FADH_2 . Molecular oxygen is required to re-oxidise the FADH_2 to FAD, with the forma-

tion of hydrogen peroxide in a 1:1 molar ratio to the resulting THCA as a by-product of the reaction (Flores-Sanchez and Verpoorte 2008, Shoyama et al. 2012). CBDA synthase also contains FAD that is reduced to FADH_2 with release of H_2O_2 , but the reaction differs from THCA synthase in the proton transfer step (Figure 10) (Taura et al. 2007a). It is estimated that about 1% of oxygen in plants is used to form reactive oxygen species in different subcellular locations with hydrogen peroxide being the most abundant. Hydrogen peroxide causes oxidative damage to cells that can lead to apoptosis (Quan et al. 2008), and plants have evolved efficient ways of eliminating toxic levels of H_2O_2 . Catalase is a peroxidase enzyme found in all oxygen-using organisms that rapidly converts H_2O_2 to water and oxygen. There are three types of catalase and the non-heme form utilises manganese (Mn^{3+}) in its catalytic center that is reduced to Mn^{2+} during the decomposition of H_2O_2 to water and oxygen. Mn^{2+} can then react with more peroxide and be converted back to Mn^{3+} according to the following equations:



Both reactions are energetically advantageous ($\Delta G < 0$). The correlation between manganese and CBN is also positive since CBN is the primary THC degradation product (Wu et al. 2004).

Bernstein et al. (2019a) describes translocation of individual macro and microelements in relation to individual plant parts' age. The work also describes, inter alia, the distribution of cannabinoids in the plant. The research shows that the concentration of cannabinoids increases with the height of the plant and the highest concentration can be found in flowers and inflorescence leaves. The concentration found in fan leaves is about 1/10 the concentration found in flowers. The distribution of mineral nutrients between plant organs shows a typical uptake and translocation in the plant. Lower concentrations of N, P, K, and higher Ca in fan leaves compared to inflorescence supports physiological findings that the fan leaves are older than the inflorescence leaves.

pH value

Suggested optimal pH range of nutrient solution is between 5.5–6.5. pH is important because it affects the availability and absorption of nutrients needed

<https://doi.org/10.17221/96/2021-PSE>




Figure 10. Reaction mechanism of tetrahydrocannabinolic acid (THCA) synthase and cannabidiolic acid (CBDA) synthase. CBGA – cannabigerolic acid; FAD – flavin adenine dinucleotide

for plant growth. In hydroponic culture, the recommended pH range is 5.5–6.0 and the maximum absorption of nutrients is usually at pH 5.8 (Velazquez et al. 2013). In growing substrate, a pH range of 5.8–7.2 is recommended and the maximum absorption of essential nutrients is typically at pH 6.5. When the pH falls below this range, many macronutrients are

less available and macronutrient deficiencies can be developed. When pH values rise above this range, many micronutrients will not be available for the plant uptake causing micronutrient deficiencies (Caplan et al. 2017a). These authors also mention the need for further research to confirm the optimal range of pH for multiple cannabis varieties.

Plant biostimulants

A plant biostimulant is any substance, micro-organism strain or mixture of both applied to plants to increase tolerance to abiotic stress, nutritional efficiency or crop quality characteristics, regardless of its nutrient content. Seven main biostimulant categories were proposed: humic and fulvic acids, protein hydrolysates, seaweed and botanical extracts, chitosan and biopolymers, beneficial bacteria, beneficial fungi and beneficial minerals (Du Jardin 2015).

Humic and fulvic acids in cannabis nutrition.

Humic substances are natural components of soil organic matter. It is a mixture of heterogeneous compounds originally classified according to their molecular weights and solubility into humins, humic acids and fulvic acids (Du Jardin 2015).

Humic acid supplementation had a positive effect on cannabis in the case of the height of cannabis plants, the chlorophyll content and the efficiency of photosynthesis, especially immediately after the period of water stress (Da Cunha Leme Filho et al. 2020).

According to the current literature, the effect on phytocannabinoids is rather negative. Bernstein et al. (2019b) mentioned that nutritional supplements such as humic acids significantly reduced spatial variability of cannabinoids throughout the plant parts. This increased uniformity came at the expenses of THC and CBD content which was reduced by 37% and 39% respectively in the top parts of plants. The decrease of THC has been associated with an additional trend of CBN increasing. This was probably due to the accelerated degradation of cannabinoids in the plant parts with their high concentration.

Other biostimulants in cannabis nutrition. Conant et al. (2017) demonstrated that microbial biostimulant Mammoth P™ promoted cannabis growth during the blooming phase. Lyu et al. (2019) hypothesised that future research will show that plant growth-promoting bacteria can affect the accumulation of phytocannabinoids, increase inflorescence yields, protect against plant pathogens by producing antimicrobial compounds and reduce the impact of abiotic stresses.

CONCLUSIONS AND FUTURE PERSPECTIVE

Based on the above information, it can be stated that quality of medical cannabis biomass, spectrum and concentration of phytocannabinoids can be influenced by cultivation conditions as well as nutrition during cultivation.

For the cultivation of medical cannabis, due to safety reasons, unpredictable environmental influences and required homogeneity of harvest, indoor cultivation is definitely a better option because optimal growing conditions can be set and cannabis can be harvested from three to six times per year. Of the growing conditions, artificial light, the level of CO₂ concentration and the humidity of the surrounding environment influence the harvest quantity and quality the most. It is very important to choose the right combination of all mentioned conditions because they affect each other.

There is currently only a few experimental data on the medical cannabis nutrition, so most of this information is based on the hemp cultivation, which was bred for fiber production rather than inflorescence. However, it can be concluded from the current literature the concentration and spectrum of individual macronutrients, micronutrients and plant biostimulants in plant nutrition has a fundamental impact on biomass formation, spectrum and amount of medical cannabis cannabinoids.

In the future, the effects of nutrient ratios and availability can reasonably be expected to be one of the main factors influencing the content and type of cannabinoids in medical cannabis plants, separate from genotype and microclimate. These issues should be explored through further experiments, which will certainly be beneficial because of growing interest in the phytocannabinoids development in public and commercial spheres. Future technical research in this area should focus on possible new indoor medical cannabis cultivation techniques or the automation of existing cultivation technologies to facilitate work.

Acknowledgement. We thank for financial support from Faculty SGS project SV20-4-21140, and prof. Gary Bentley for improving the use of English in the manuscript.

REFERENCES

- Amaducci S., Scordia D., Liu F., Zhang Q., Guo H., Testa G., Cosentino S. (2015): Key cultivation techniques for hemp in Europe and China. *Industrial Crops and Products*, 68: 2–16.
- Aubin M.-P., Seguin P., Vanasse A., Tremblay G.F., Mustafa A.F., Charron J.-B. (2015): Industrial hemp response to nitrogen, phosphorus, and potassium fertilization. *Crop, Forage and Turfgrass Management*, 1: 1–10.
- Babson K.A., Sottile J., Morabito D. (2017): Cannabis, cannabinoids, and sleep: a review of the literature. *Current Psychiatry Reports*, 19: 1–12.

<https://doi.org/10.17221/96/2021-PSE>

- Balneaves L.G., Alraja A.A. (2019): "Guarding their practice": a descriptive study of Canadian nursing policies and education related to medical cannabis. *BMC Nursing*, 18: 1–10.
- Barker A.V., Pilbeam D.J. (2015): *Handbook of Plant Nutrition*. Boca Raton, CRC Press. ISBN 9781032098630
- Bazzaz F., Dusek D., Seigler D., Haney A. (1975): Photosynthesis and cannabinoid content of temperate and tropical populations of *Cannabis sativa*. *Biochemical Systematics and Ecology*, 3: 15–18.
- Bernstein N., Gorelick J., Koch S. (2019a): Interplay between chemistry and morphology in medical cannabis (*Cannabis sativa* L.). *Industrial Crops and Products*, 129: 185–194.
- Bernstein N., Gorelick J., Zerahia R., Koch S. (2019b): Impact of N, P, K and humic acids supplementation on the chemical profile of medical cannabis (*Cannabis sativa* L.). *Frontiers in Plant Science*, 10: 736.
- Bessho M., Shimizu K. (2012): Latest trends in LED lighting. *Electronics and Communications in Japan*, 95: 1–7.
- Beutler J.A., Marderosian A.H. (1978): Chemotaxonomy of Cannabis I. Crossbreeding between *Cannabis sativa* and *C. ruderalis*, with analysis of cannabinoid content. *Economic Botany*, 32: 387–394.
- Bócsa I., Máthé P., Hangyel L. (1997): Effect of nitrogen on tetrahydrocannabinol (THC) content in hemp (*Cannabis sativa* L.) leaves at different positions. *Journal of the International Hemp Association*, 4: 78–79.
- Bonn-Miller M.O., Babson K.A., Vandrey R. (2014): Using cannabis to help you sleep: heightened frequency of medical cannabis use among those with PTSD. *Drug and Alcohol Dependence*, 136: 162–165.
- Boon E.M., Downs A., Marcey D. (2007): Catalase: $H_2O_2:H_2O_2$ Oxidoreductase. *Catalase Structural Tutorial Text*, 2007–2002.
- Brickell C., Alexander C., David J., Hetterscheid W., Leslie A., Malécot V., Xiaobai J. (2009): *International Code of Nomenclature for Cultivated Plants*. Leuven, International Society for Horticultural Science.
- Caplan D., Dixon M., Zheng Y. (2017a): Optimal rate of organic fertilizer during the flowering stage for cannabis grown in two coir-based substrates. *HortScience*, 52: 1796–1803.
- Caplan D., Dixon M., Zheng Y. (2017b): Optimal rate of organic fertilizer during the vegetative-stage for cannabis grown in two coir-based substrates. *HortScience*, 52: 1307–1312.
- Carpentier C., Mulligan K., Laniel L., Potter D., Hughes B., Vandam L., Skarupova K. (2012): *Cannabis Production and Markets in Europe*. Lisbon, Publications Office of the European Union. ISBN 978-92-9168-502-8
- Coffman C., Gentner W. (1975): Cannabinoid profile and elemental uptake of *Cannabis sativa* L. as influenced by soil characteristics. *Agronomy Journal*, 67: 491–497.
- Coffman C., Gentner W. (1977): Responses of greenhouse-grown *Cannabis sativa* L. to nitrogen, phosphorus, and potassium. *Agronomy Journal*, 69: 832–836.
- Conant R., Walsh R., Walsh M., Bell C., Wallenstein M. (2017): Effects of a microbial biostimulant, Mammoth PTM, on *Cannabis sativa* bud yield. *Journal of Horticulture*, 4: 1000191.
- Corey-Bloom J., Wolfson T., Gamst A., Jin S., Marcotte T.D., Bentley H., Gouaux B. (2012): Smoked cannabis for spasticity in multiple sclerosis: a randomized, placebo-controlled trial. *Canadian Medical Association Journal*, 184: 1143–1150.
- Croteau R. (1987): Biosynthesis and catabolism of monoterpenoids. *Chemical Reviews*, 87: 929–954.
- Da Cunha Leme Filho J.F., Thomason W.E., Evanylo G.K., Zhang X., Strickland M.S., Chim B.K., Diatta A.A. (2020): Biochemical and physiological responses of *Cannabis sativa* to an integrated plant nutrition system. *Agronomy Journal*, 112: 5237–5248.
- De Backer B., Maebe K., Verstraete A.G., Charlier C. (2012): Evolution of the content of THC and other major cannabinoids in drug-type cannabis cuttings and seedlings during growth of plants. *Journal of Forensic Sciences*, 57: 918–922.
- De Lamarck J.-B.D.M. (1786): *Methodical Encyclopedia: Botany*. Paris, Panckoucke. (In French)
- Devane W.A., Hanus L., Breuer A., Pertwee R.G., Stevenson L.A., Griffin G., Gibson D., Mandelbaum A., Etinger A., Mechoulam R. (1992): Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science*, 258: 1946–1949.
- Dewick P.M. (2002): *Medicinal Natural Products: A Biosynthetic Approach*. Chichester, John Wiley and Sons. ISBN: 978-0-470-74168-9
- Doyle E., Spence A. (1995): Cannabis as a medicine? *British Journal of Anaesthesia*, 74: 359–361.
- Drugs U.N.O.O. (2009): *Recommended Methods for the Identification and Analysis of Cannabis and Cannabis Products: Manual for Use by National Drug Testing Laboratories*. Vienna, United Nations Publications. ISBN: 9789210555890
- Du Jardin P. (2015): Plant biostimulants: definition, concept, main categories and regulation. *Scientia Horticulturae*, 196: 3–14.
- Duran M., Pérez E., Abanades S., Vidal X., Saura C., Majem M., Arriola E., Rabanal M., Pastor A., Farré M. (2010): Preliminary efficacy and safety of an oromucosal standardized cannabis extract in chemotherapy-induced nausea and vomiting. *British Journal of Clinical Pharmacology*, 70: 656–663.
- Elsehly M.A., Slade D. (2005): Chemical constituents of marijuana: the complex mixture of natural cannabinoids. *Life Sciences*, 78: 539–548.
- Elzinga S., Fishedick J., Podkolinski R., Raber J. (2015): Cannabinoids and terpenes as chemotaxonomic markers in cannabis. *Natural Products Chemistry and Research*, 3: 181.
- Evans J., Jakobsen I., Ögren E. (1993): Photosynthetic light-response curves. *Planta*, 189: 191–200.
- Farag S., Kayser O. (2015): Cultivation and breeding of *Cannabis sativa* L. for preparation of standardized extracts for medicinal purposes. *Medicinal and Aromatic Plants of the World*. Medicinal and Aromatic Plants of the World, 165–186.
- Fellermeier M., Zenk M.H. (1998): Prenylation of olivetolate by a hemp transferase yields cannabigerolic acid, the precursor of tetrahydrocannabinol. *FEBS Letters*, 427: 283–285.
- Fishedick J.T., Hazekamp A., Erkelens T., Choi Y.H., Verpoorte R. (2010): Metabolic fingerprinting of *Cannabis sativa* L., cannabi-

<https://doi.org/10.17221/96/2021-PSE>

- noids and terpenoids for chemotaxonomic and drug standardization purposes. *Phytochemistry*, 71: 2058–2073.
- Flores-Sanchez I.J., Verpoorte R. (2008): PKS activities and biosynthesis of cannabinoids and flavonoids in *Cannabis sativa* L. plants. *Plant and Cell Physiology*, 49: 1767–1782.
- Gagne S.J., Stout J.M., Liu E., Boubakir Z., Clark S.M., Page J.E. (2012): Identification of olivetolic acid cyclase from *Cannabis sativa* reveals a unique catalytic route to plant polyketides. *Proceedings of the National Academy of Sciences of the United States of America*, 109: 12811–12816.
- Gaoni Y., Mechoulam R. (1964): Isolation, structure, and partial synthesis of an active constituent of hashish. *Journal of the American Chemical Society*, 86: 1646–1647.
- Giuffrida A., Beltramo M., Piomelli D. (2001): Mechanisms of endocannabinoid inactivation: biochemistry and pharmacology. *Journal of Pharmacology and Experimental Therapeutics*, 298: 7–14.
- Grotenhermen F., Muller-Vahl K. (2012): The therapeutic potential of cannabis and cannabinoids. *Deutsches Aertzblatt International*, 109: 495–501.
- Halliday K.J., Fankhauser C. (2003): Phytochrome-hormonal signalling networks. *New Phytologist*, 157: 449–463.
- Hamilton E.W., Heckathorn S.A., Joshi P., Wang D., Barua D. (2008): Interactive effects of elevated CO₂ and growth temperature on the tolerance of photosynthesis to acute heat stress in C3 and C4 species. *Journal of Integrative Plant Biology*, 50: 1375–1387.
- Haney A., Kutscheid B.B. (1973): Quantitative variation in the chemical constituents of marihuana from stands of naturalized *Cannabis sativa* L. in East-Central Illinois. *Economic Botany*, 27: 193–203.
- Hanuš L., Dostálová M. (1994): The effect of soil fertilization on the formation and the amount of cannabinoid substances in *Cannabis sativa* L. in the course of one vegetation period. *Acta Universitatis Palackianae Olomucensis Facultatis Medicae*, 138: 11–15.
- Hanuš L.O., Meyer S.M., Muñoz E., Tagliabata Scafati O., Appendino G. (2016): Phytocannabinoids: a unified critical inventory. *Natural Product Reports*, 33: 1357–1392.
- Hartsel J.A., Eades J., Hickory B., Makriyannis A. (2016): *Cannabis sativa* and Hemp. *Nutraceuticals*. Elsevier, 735–754.
- Hemphill J.K., Turner J.C., Mahlberg P.G. (1980): Cannabinoid content of individual plant organs from different geographical strains of *Cannabis sativa* L. *Journal of Natural Products*, 43: 112–122.
- Hillig K.W. (2004): A chemotaxonomic analysis of terpenoid variation in Cannabis. *Biochemical Systematics and Ecology*, 32: 875–891.
- Hillig K.W. (2005): Genetic evidence for speciation in Cannabis (Cannabaceae). *Genetic Resources and Crop Evolution*, 52: 161–180.
- Hillig K.W., Mahlberg P.G. (2004): A chemotaxonomic analysis of cannabinoid variation in Cannabis (Cannabaceae). *American Journal of Botany*, 91: 966–975.
- Chandra S., Lata H., Khan I.A., Elsohly M.A. (2008): Photosynthetic response of *Cannabis sativa* L. to variations in photosynthetic photon flux densities, temperature and CO₂ conditions. *Physiology and Molecular Biology of Plants*, 14: 299–306.
- Chandra S., Lata H., Khan I.A., Elsohly M.A. (2011): Photosynthetic response of *Cannabis sativa* L., an important medicinal plant, to elevated levels of CO₂. *Physiology and Molecular Biology of Plants*, 17: 291–295.
- Janischevsky D. (1924): A form of hemp in wild areas of southeastern Russia. *NG Černyševskogo Universiteta*, 2: 3–17.
- Järvinen T., Pate D.W., Laine K. (2002): Cannabinoids in the treatment of glaucoma. *Pharmacology and Therapeutics*, 95: 203–220.
- Jiang H.-E., Li X., Zhao Y.-X., Ferguson D.K., Hueber F., Bera S., Wang Y.-F., Zhao L.-C., Liu C.-J., Li C.-S. (2006): A new insight into *Cannabis sativa* (Cannabaceae) utilization from 2 500-year-old Yanghai Tombs, Xinjiang, China. *Journal of Ethnopharmacology*, 108: 414–422.
- Kimball B.A. (1983): Carbon dioxide and agricultural yield: an assemblage and analysis of 430 prior observations 1. *Agronomy Journal*, 75: 779–788.
- Kimura M., Okamoto K. (1970): Distribution of tetrahydrocannabinolic acid in fresh wild cannabis. *Experientia*, 26: 819–820.
- Kriesche U., Schumann E., Weber W., Beyer M., Brühl L. (2004): Oil content, tocopherol composition and fatty acid patterns of the seeds of 51 *Cannabis sativa* L. genotypes. *Euphytica*, 137: 339–351.
- Lane M., Vogel C.L., Ferguson J., Krasnow S., Saiers J.L., Hamm J., Salva K., Wiernik P.H., Holroyde C.P., Hammill S. (1991): Dronabinol and prochlorperazine in combination for treatment of cancer chemotherapy-induced nausea and vomiting. *Journal of Pain and Symptom Management*, 6: 352–359.
- Leggett T. (2006): A review of the world cannabis situation. *Bulletin on Narcotics*, 58: 1–155.
- Linnaeus C. (1753): *Species Plantarum*. Laurentius Salvius, Sweden.
- Lotan I., Treves T.A., Roditi Y., Djaldetti R. (2014): Cannabis (medical marijuana) treatment for motor and non-motor symptoms of Parkinson disease: an open-label observational study. *Clinical Neuropharmacology*, 37: 41–44.
- Lydon J., Teramura A.H., Coffman C.B. (1987): UV-B radiation effects on photosynthesis, growth and cannabinoid production of two *Cannabis sativa* chemotypes. *Photochemistry and Photobiology*, 46: 201–206.
- Lynch M.E., Campbell F. (2011): Cannabinoids for treatment of chronic non-cancer pain; a systematic review of randomized trials. *British Journal of Clinical Pharmacology*, 72: 735–744.
- Lyu D., Backer R., Robinson W.G., Smith D.L. (2019): Plant growth-promoting rhizobacteria for cannabis production: yield, cannabinoid profile and disease resistance. *Frontiers in Microbiology*, 10: 1761.
- Magagnoli G., Grassi G., Kotiranta S. (2018): The effect of light spectrum on the morphology and cannabinoid content of *Cannabis sativa* L. *Medical Cannabis and Cannabinoids*, 1: 19–27.
- Mahlberg P.G., Hemphill J.K. (1983): Effect of light quality on cannabinoid content of *Cannabis sativa* L. (Cannabaceae). *Botanical Gazette*, 144: 43–48.
- Massa G.D., Kim H.-H., Wheeler R.M., Mitchell C.A. (2008): Plant productivity in response to LED lighting. *HortScience*, 43: 1951–1956.

<https://doi.org/10.17221/96/2021-PSE>

- McNeill J., Barrie F., Buck W., Demoulin V., Greuter W., Hawksworth D., Herendeen P., Knapp S., Marhold K., Prado J. (2012): International Code of Nomenclature for Algae, Fungi and Plants (Melbourne Code). Koeltz Scientific Books Königstein.
- McPartland J.M. (2018): Cannabis systematics at the levels of family, genus, and species. *Cannabis and Cannabinoid Research*, 3: 203–212.
- Mechoulam R., Ben-Shabat S., Hanus L., Ligumsky M., Kaminski N.E., Schatz A.R., Gopher A., Almog S., Martin B.R., Compton D.R., Pertwee R.G., Griffin G., Bayewitch M., Barg J., Vogel Z. (1995): Identification of an endogenous 2-monoglyceride, present in canine gut, that binds to cannabinoid receptors. *Biochemical Pharmacology*, 50: 83–90.
- Mechoulam R., Braun P., Gaoni Y. (1967): A stereospecific synthesis of (-)-delta 1- and (-)-delta 1(6)-tetrahydrocannabinols. *Journal of the American Chemical Society*, 89: 4552–4554.
- Mechoulam R., Gaoni Y. (1967): The absolute configuration of delta-1-tetrahydrocannabinol, the major active constituent of hashish. *Tetrahedron Letters*, 12: 1109–1111.
- Milay L., Berman P., Shapira A., Guberman O., Meiri D. (2020): Metabolic profiling of cannabis secondary metabolites for evaluation of optimal postharvest storage conditions. *Frontiers in Plant Science*, 11: 583605.
- Minorsky P.V. (2002): Global Warming – Effects on Plants. *Plant Physiology*, 129: 1421–1422.
- Mölleken H., Theimer R.R. (1997): Survey of minor fatty acids in *Cannabis sativa* L. fruits of various origins. *Journal of the International Hemp Association*, 4: 13–18.
- Morimoto S., Komatsu K., Taura F., Shoyama Y. (1997): Enzymological evidence for cannabichromenic acid biosynthesis. *Journal of Natural Products*, 60: 854–857.
- Morimoto S., Komatsu K., Taura F., Shoyama Y. (1998): Purification and characterization of cannabichromenic acid synthase from *Cannabis sativa*. *Phytochemistry*, 49: 1525–1529.
- Namdar D., Charuvi D., Ajampura V., Mazuz M., Ion A., Kamara I., Koltai H. (2019): LED lighting affects the composition and biological activity of *Cannabis sativa* secondary metabolites. *Industrial Crops and Products*, 132: 177–185.
- Ögren E., Evans J. (1993): Photosynthetic light-response curves. *Planta*, 189: 182–190.
- Pate D.W. (1994): Chemical ecology of Cannabis. *Journal of the International Hemp Association*, 2: 32–37.
- Patel S., Cudney R., McPherson A. (1994): Crystallographic characterization and molecular symmetry of edestin, a legumin from hemp. *Journal of Molecular Biology*, 235: 361–363.
- Pickering K.L., Beckermann G., Alam S., Foreman N.J. (2007): Optimising industrial hemp fibre for composites. *Composites Part A: Applied Science and Manufacturing*, 38: 461–468.
- Pooyania S., Ethans K., Szturm T., Casey A., Perry D. (2010): A randomized, double-blinded, crossover pilot study assessing the effect of nabilone on spasticity in persons with spinal cord injury. *Archives of Physical Medicine and Rehabilitation*, 91: 703–707.
- Portenoy R.K., Ganae-Motan E.D., Allende S., Yanagihara R., Shaiova L., Weinstein S., McQuade R., Wright S., Fallon M.T. (2012): Nabiximols for opioid-treated cancer patients with poorly-controlled chronic pain: a randomized, placebo-controlled, graded-dose trial. *The Journal of Pain*, 13: 438–449.
- Potter D.J. (2014): A review of the cultivation and processing of cannabis (*Cannabis sativa* L.) for production of prescription medicines in the UK. *Drug Testing and Analysis*, 6: 31–38.
- Potter D.J., Duncombe P. (2012): The effect of electrical lighting power and irradiance on indoor-grown cannabis potency and yield. *Journal of Forensic Sciences*, 57: 618–622.
- Quan L.J., Zhang B., Shi W.W., Li H.Y. (2008): Hydrogen peroxide in plants: a versatile molecule of the reactive oxygen species network. *Journal of Integrative Plant Biology*, 50: 2–18.
- Radosavljevic-Stevanovic N., Markovic J., Agatonovic-Kustrin S., Razic S. (2014): Metals and organic compounds in the biosynthesis of cannabinoids: a chemometric approach to the analysis of *Cannabis sativa* samples. *Natural Product Research*, 28: 511–516.
- Ravikoff Allegratti J., Courtwright A., Lucci M., Korzenik J.R., Levine J. (2013): Marijuana use patterns among patients with inflammatory bowel disease. *Inflammatory Bowel Diseases*, 19: 2809–2814.
- Razdan R.K. (1986): Structure-activity relationships in cannabinoids. *Pharmacological Reviews*, 38: 75–149.
- Ross S., Elsohly M. (1995): Constituents of *Cannabis sativa* L. XXVIII – A review of the natural constituents: 1980–1994. *Zagazig Journal of Pharmaceutical Sciences*, 4: 1–10.
- Russo E.B., Guy G.W., Robson P.J. (2007): Cannabis, pain, and sleep: lessons from therapeutic clinical trials of Sativex®, a cannabis-based medicine. *Chemistry and Biodiversity*, 4: 1729–1743.
- Saloner A., Sacks M.M., Bernstein N. (2019): Response of medical cannabis (*Cannabis sativa* L.) genotypes to K supply under long photoperiod. *Frontiers in Plant Science*, 10: 1369.
- Selvarajah D., Gandhi R., Emery C.J., Tesfaye S. (2010): Randomized placebo-controlled double-blind clinical trial of cannabis-based medicinal product (Sativex) in painful diabetic neuropathy: depression is a major confounding factor. *Diabetes Care*, 33: 128–130.
- Shelef A., Mashiah M., Schumacher I., Shine O., Baruch Y. (2011): Medical grade cannabis (MGC): regulation mechanisms, the present situation around the world and in Israel. *Harefuah*, 150: 913–917, 935, 934.
- Shoyama Y., Hirano H., Nishioka I. (1984): Biosynthesis of propyl cannabinoid acid and its biosynthetic relationship with pentyl and methyl cannabinoid acids. *Phytochemistry*, 23: 1909–1912.
- Shoyama Y., Tamada T., Kurihara K., Takeuchi A., Taura F., Arai S., Blaber M., Shoyama Y., Morimoto S., Kuroki R. (2012): Structure and function of Δ^1 -tetrahydrocannabinolic acid (THCA) synthase, the enzyme controlling the psychoactivity of *Cannabis sativa*. *Journal of Molecular Biology*, 423: 96–105.
- Shoyama Y., Yamauchi T., Nishioka I. (1970): Cannabis. V. Cannabigerolic acid monomethyl ether and cannabinolic acid. *Chemical and Pharmaceutical Bulletin*, 18: 1327–1332.

- Small E. (2015): Evolution and classification of *Cannabis sativa* (marijuana, hemp) in relation to human utilization. *The Botanical Review*, 81: 189–294.
- Small E., Cronquist A. (1976): A practical and natural taxonomy for Cannabis. *Taxon*, 25: 405–435.
- Surragé V.A., Lafrenière C., Dixon M., Zheng Y. (2010): Benefits of vermicompost as a constituent of growing substrates used in the production of organic greenhouse tomatoes. *HortScience*, 45: 1510–1515.
- Sweet S.L. (2016): The energy intensity of lighting used for the production of recreational cannabis in Washington State and implications for energy efficiency. Olympia, Evergreen State College.
- Taura F. (2009): Studies on tetrahydrocannabinolic acid synthase that produces the acidic precursor of tetrahydrocannabinol, the pharmacologically active cannabinoid in marijuana. *Drug Discoveries and Therapeutics*, 3: 83–87.
- Taura F., Morimoto S., Shoyama Y. (1995a): Cannabinerolic acid, a cannabinoid from *Cannabis sativa*. *Phytochemistry*, 39: 457–458.
- Taura F., Morimoto S., Shoyama Y. (1996): Purification and characterization of cannabidiolic-acid synthase from *Cannabis sativa* L. Biochemical analysis of a novel enzyme that catalyzes the oxidocyclization of cannabigerolic acid to cannabidiolic acid. *Journal of Biological Chemistry*, 271: 17411–17416.
- Taura F., Morimoto S., Shoyama Y., Mechoulam R. (1995b): First direct evidence for the mechanism of Δ^9 -THC biosynthesis. *Journal of the American Chemical Society*, 117: 9766–9767.
- Taura F., Sirikantaramas S., Shoyama Y., Shoyama Y., Morimoto S. (2007a): Phytocannabinoids in *Cannabis sativa*: recent studies on biosynthetic enzymes. *Chemistry and Biodiversity*, 4: 1649–1663.
- Taura F., Sirikantaramas S., Shoyama Y., Yoshikai K., Shoyama Y., Morimoto S. (2007b): Cannabidiolic-acid synthase, the chemotype-determining enzyme in the fiber-type *Cannabis sativa*. *FEBS letters*, 581: 2929–2934.
- Taura F., Tanaka S., Taguchi C., Fukamizu T., Tanaka H., Shoyama Y., Morimoto S. (2009): Characterization of olivetol synthase, a polyketide synthase putatively involved in cannabinoid biosynthetic pathway. *FEBS Letters*, 583: 2061–2066.
- Toonen M., Ribot S., Thissen J. (2006): Yield of illicit indoor cannabis cultivation in the Netherlands. *Journal of Forensic Sciences*, 51: 1050–1054.
- Troutt W.D., Didonato M.D. (2015): Medical cannabis in Arizona: patient characteristics, perceptions, and impressions of medical cannabis legalization. *Journal of Psychoactive Drugs*, 47: 259–266.
- Turner C.E., Elsohly M.A., Boeren E.G. (1980): Constituents of *Cannabis sativa* L. XVII. A review of the natural constituents. *Journal of Natural Products*, 43: 169–234.
- Van Bakel H., Stout J.M., Cote A.G., Tallon C.M., Sharpe A.G., Hughes T.R., Page J.E. (2011): The draft genome and transcriptome of *Cannabis sativa*. *Genome Biology*, 12: 1–18.
- Vanhove W., Surmont T., Van Damme P., De Ruyver B. (2012): Yield and turnover of illicit indoor cannabis (*Cannabis* spp.) plantations in Belgium. *Forensic Science International*, 220: 265–270.
- Vanhove W., Van Damme P., Meert N. (2011): Factors determining yield and quality of illicit indoor cannabis (*Cannabis* spp.) production. *Forensic Science International*, 212: 158–163.
- Velazquez L., Hernandez M., Leon M., Domínguez R.B., Gutierrez J. (2013): First advances on the development of a hydroponic system for cherry tomato culture. 10th International Conference on Electrical Engineering, Computing Science and Automatic Control (CCE). IEEE.
- Vera C., Malhi S., Raney J., Wang Z. (2004): The effect of N and P fertilization on growth, seed yield and quality of industrial hemp in the Parkland region of Saskatchewan. *Canadian Journal of Plant Science*, 84: 939–947.
- Vlasits J., Jakopitsch C., Bernroither M., Zamocky M., Furtmüller P.G., Obinger C. (2010): Mechanisms of catalase activity of heme peroxidases. *Archives of Biochemistry and Biophysics*, 500: 74–81.
- Wade D.T., Makela P., Robson P., House H., Bateman C. (2004): Do cannabis-based medicinal extracts have general or specific effects on symptoms in multiple sclerosis? A double-blind, randomized, placebo-controlled study on 160 patients. *Multiple Sclerosis Journal*, 10: 434–441.
- Wang D., Heckathorn S.A., Barua D., Joshi P., Hamilton E.W., Lacroix J.J. (2008): Effects of elevated CO₂ on the tolerance of photosynthesis to acute heat stress in C₃, C₄, and CAM species. *American Journal of Botany*, 95: 165–176.
- Wheeler T., Batts G., Ellis R., Hadley P., Morison J. (1996): Growth and yield of winter wheat (*Triticum aestivum*) crops in response to CO₂ and temperature. *The Journal of Agricultural Science*, 127: 37–48.
- Wilsey B., Marcotte T., Deutsch R., Gouaux B., Sakai S., Donaghe H. (2013): Low-dose vaporized cannabis significantly improves neuropathic pain. *The Journal of Pain*, 14: 136–148.
- Wu A.J., Penner-Hahn J.E., Pecoraro V.L. (2004): Structural, spectroscopic, and reactivity models for the manganese catalases. *Chemical Reviews*, 104: 903–938.
- Yu Y. (1987): Agricultural history over seven thousand years in China. In: Wittwer S., Yu Y.T., Han S., Wang L.Z. (eds.): *Feeding a Billion – Frontiers of Chinese Agriculture*. East Lansing, Michigan State University Press, 19–33.
- Zelitch I. (1975): Improving the efficiency of photosynthesis. *Science*, 188: 626–633.
- Zhai Z., Ehret D.L., Forge T., Helmer T., Lin W., Dorais M., Papadopoulos A.P. (2009): Organic fertilizers for greenhouse tomatoes: productivity and substrate microbiology. *HortScience*, 44: 800–809.
- Zhang Q., Chen X., Guo H., Trindade L.M., Salentijn E.M., Guo R., Guo M., Xu Y., Yang M. (2018): Latitudinal adaptation and genetic insights into the origins of *Cannabis sativa* L. *Frontiers in Plant Science*, 9: 1876.
- Zlas J., Stark H., Seligman J., Levy R., Werker E., Breuer A., Mechoulam R. (1993): Early medical use of cannabis. *Nature*, 363: 215–215.
- Zuk-Golaszewska K., Golaszewski J. (2018): *Cannabis sativa* L. – cultivation and quality of raw material. *Journal of Elementology*, 23: 971–984.

Received: February 23, 2021

Accepted: June 4, 2021

Published online: July 19, 2021